



FACT SHEET

FABRIC PESTS

The group of insects known as fabric pests includes silverfish, clothes moths and carpet beetles. Except for the silverfish, the main food constituent sought after by fabric pests, is a protein substance called Keratin present in animal fibers such as wool. Silverfish prefer to eat substances of plant origin such as cotton cellulose and starch.

SILVER FISH



Appearance: Silverfish (*Ctenolepisma longicaudata*) are wingless, primitive insects and can easily be identified by the three long appendages protruding from the rear of the abdomen. They have a fish-like appearance with the body tapering to the rear and are covered in scales giving them a dull silvery colouration.

Lifecycle: The female adult lays eggs either singly or in small batches of (2-20eggs) which usually hatch after a period of 2 – 8 weeks into nymphs which closely resemble the shape of the adult but are smaller in size. The nymphs undergo a series of moults over a period of 3 – 24 months becoming sexually mature adults. They are one of the few insects to continue moulting after they have achieved adulthood and can go on to live for a period of up to 4 years and a full year without eating.

Behavior & Habits: Silverfish can move extremely quickly when disturbed and in general tend to avoid light. They are usually found in dark undisturbed areas; however, they often range throughout a building, particularly in cupboards, stored paper and bookshelves and behind wallpaper that has peeled away from the wall surface. Silverfish cannot climb smooth surfaces and so are often found trapped in glasses, baths, and basin. Silverfish do not carry disease organisms harmful to man or domestic animals.

Preventative & Control: Is normally achieved by the storage of books, paper and linen fabrics in well lit and ventilated areas. Chemical control consists of the careful application of residual surface sprays and the judicious use of space sprays. The client will need to empty all cupboards including kitchen and pantry, wardrobes, linen and vanity cupboards so Exopest can apply a registered pesticide to both topside and underside of each shelf. This is necessary as silverfish live in cracks and crevices. This preparation must occur prior to our serviceman arriving. The use of insecticidal dust into cracks and crevices around architraves, skirting's, windows and doors and other likely harborages will also be done. Roof and subfloor also to be treated

CARPET BEETLES

The most common carpet beetles to be found attacking various fabrics of animal origin are the black carpet beetle (*Attagenus unicolor*) and the variegated carpet beetle (*Anthrenus verbasci*). As well as carpets, they can be found in woolen goods, fur, silk, upholstery and stuffed animals.



An infestation of carpet beetles usually occurs in undisturbed areas, such as below items of heavy furniture or perimeter areas of the carpet. It is possible for the carpet beetles to attack synthetic carpets particularly if they are heavily soiled with organic food debris or urine. Many synthetic and woolen carpets are now treated with insecticides at the manufacturing stage rendering them resistant to carpet beetle attacks.

Habits & Lifecycle: The adult female beetle lays her eggs in dark undisturbed areas. The eggs hatch into reddy-brown colored larvae or grubs covered with tufted hairs. It is this larval stage that causes the damage to fabrics, rugs, underfelts and carpets. When they are fully fed the larvae change into immobile pupae stage, that after several weeks turn into adult beetles. The adult beetles usually fly to exterior areas where they are often attracted to white flowers. They cause no damage but often infest dwellings by their presence in birds' nests or being introduced to the interior of dwellings on cut flowers.

Control of Carpet Beetle: is usually achieved by attention to thorough vacuuming of the infested areas followed Exopest applying a residual insecticide sprays. The client will need to empty cupboards including, wardrobes, and linen cupboards so application to both topside and underside of each shelf can occur. This preparation must occur prior to our serviceman arriving. The use of insecticidal dust into cracks and crevices around architraves, skirting's, windows and doors and other likely harbourages will also be done. The roof and subfloor may need to be treated, usually with a dust formulation.

CLOTHES MOTHS



Although less common than carpet beetles, these fabric pests are commonly experienced in similar environments to carpet beetles, namely woolen materials, felt, fur and other materials of animal origin. The two most common species of clothes moths are the case making clothes moth (*Tinea pellionella*) and the common clothes moth (*Tineola bisselliella*).

Habits & Lifecycle

The female clothes moths lay eggs on materials that will provide an adequate food source for the emerging larvae. The larvae or grubs feed in dark undisturbed areas and larvae can be protected by a "case" woven from the fibers of the material on which they are feeding, (case making clothes moth) or they may feed in tubes constructed in the material being attacked (common clothes moth). When the larvae are fully fed the case making clothes moth larvae migrate to areas high up on the wall or on the ceiling to pupate, whereas the common clothes moth pupates within the feeding material.

The pupa of both species change into the adult moth and are usually short lived. Signs of clothes moth activity can be indicated by the presence of damaged material, cast larval skins and sand like larval droppings.

Control of Clothes Moth: generally consists of Exopest applying low toxic residual preparations to affected areas. It may be necessary to space spray affected areas to control adult Clothes Moths. **The same preparation applies as Carpet Beetle for the client.** When storing clothes for long periods, these should be dry cleaned before being stored in clean plastic bags with moth balls.

You may experience several adult moth hatchings initially 2-3 weeks after treatment, and again several months later, some will be caught in the pheromone traps, others you can just spray with an aerosol. Treatment does not kill the eggs but does kill the moth larvae after they hatch out and crawl along treated surfaces. Depending on when the eggs were laid you may see live larvae hatch out several months after treatment. This is quite normal and may see them crawl out into open areas, away from the treated edge and die off.

